Fluid Phase Equilibria, 39 (1988) 335–336
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands

Contents Volume 39 (1988)

Volume 39, No. 1

Generalized equation of state for Lennard–Jones fluids—I. Pure fluids and simple mixtures
Y. Adachi (Nagoya, Japan), I. Fijihara (Osaka, Japan), M. Takamiya and K. Nakanishi (Aichi, Japan) 1–38

Thermodynamics of associating component + saturated hydrocarbon mixtures at low pressures—IV. Correlation of vapour pressures and volumetric properties of some aliphatic amines and their mixtures with n-alkanes in terms of association
I. Cibulka, K. Tamura and I. Nagata (Kanazawa, Japan) 39–51

Calculation of tri-critical points
M.L. Michelsen (Lyngby, Denmark) and R.A. Heidemann (Calgary, Alb., Canada) 53–74

A simple and accurate technique to obtain pure component parameters for three-parameter equations of state
L.A. Cisternas (Antofagasta, Chile) 75–87

Application of hard convex body and hard sphere equations of state to the critical properties of binary mixtures
R.J. Sadus, C.L. Young (Parkville, Vic., Australia) and P. Svejda (Bochum, F.R.G.) 89–99

Bubble-point pressures of the H_2–CO–CO_2 system
C. Yokoyama, K. Arai, S. Saito (Sendai, Japan) and H. Mori (Tokyo, Japan) . 101–110

Volume 39, No. 2

Thermodynamic property predictions with the Trebble–Bishnoi equation of state
M.A. Trebble and P.R. Bishnoi (Calgary, Alberta, Canada) 111–128

Extension of the Associated Perturbed Anisotropic Chain Theory to mixtures with more than one associating component
G.D. Ikonomou and M.D. Donohue (Baltimore, MD, U.S.A.) 129–159

Application of the Kirkwood–Buff theory of solutions to dilute supercritical mixtures. II. The excluded volume and local composition models
D.M. Pfund, L.L. Lee (Norma, OK, U.S.A.) and H.D. Cochran (Oak Ridge, TN, U.S.A.) 161–192

Thermodynamic properties of ethanol and water—III. Description of the different excess functions (gE, hE, c_p^E) using an empirical model
A. Gaulhofer, B. Kolbe and J. Gmehling (Dortmund, F.R.G.) 193–209

High-pressure phase equilibria in the system glucose + fructose + water + ethanol + carbon dioxide
R. D'Souza and A.S. Teja (Atlanta, GA, U.S.A.) 211–224

Erratum 225

0378-3812/88/$03.50 © 1988 Elsevier Science Publishers B.V.

Volume 39, No. 3

An analytical model for aqueous electrolyte solutions based on fluctuation solution theory
R.L. Perry, J.D. Massie and P.T. Cummings (Charlottesville, VA, U.S.A.) . . . 227–266
Heats of mixing-composition isotherms deduced from only two ($g^E - x$) isotherms
D.A. Jonah (Sierra Leone, West Africa) 267–291
Vapour–liquid equilibria of dilute solutions of phenols in cyclohexane. Self-association of *ortho*-substituted phenols
R. Bareła and H. Buchowski (Warsaw, Poland) 293–306
Measurements of VLE, v^E and h^E for binary mixtures of n-alkanes with chloro-alkylbenzenes
H.I. Paul, J. Krug and H. Knapp (Berlin, F.R.G.) 307–324

Short Communication

The mapping of three phase volumetric behavior of pseudo-binary systems on pressure-composition diagrams
N. Pollack and R. Enick (Pittsburg, PA, U.S.A.) 325–332

Author Index 333

Contents of Volume 39 (1988) 335–336